

Letters to the Editor

Stroke welcomes Letters to the Editor and will publish them, if suitable, as space permits. They should not exceed 750 words (including references) and may be subject to editing or abridgment. Please submit letters in duplicate, typed double-spaced. Include a fax number for the corresponding author and a completed copyright transfer agreement form (published in every issue).

Multifactorial Bases of Stroke After Cardiac Surgery

To the Editor:

We read with great interest the article by Gottesman et al¹ dealing with one of the most devastating complications of cardiac surgery, perioperative stroke. The article analyzed the impact on outcomes of watershed stroke and the role of intraoperative hypoperfusion on its onset. Nevertheless, their report raises some concerns.

The authors pointed out that hypoperfusion is an independent predictor of bilateral watershed strokes, a drop in mean arterial pressure of at least 10 mm Hg being associated with a 4-time increased adjusted risk of watershed strokes. The study group was composed of patients who developed postoperative stroke whereas patients who underwent cardiac surgery without developing stroke were not considered for statistical analysis. Moreover, only the relationship between bilateral watershed strokes and hypoperfusion was analyzed, whereas monolateral watershed strokes were considered together with other stroke patterns in group 2, despite the underlying mechanisms being more similar in monolateral and bilateral stroke than in other stroke patterns. Hence, by its nature this interesting study evaluated the impact of hypoperfusion on stroke severity whereas no conclusion can be given on the role of changes in blood pressure on stroke incidence and stroke patterns.

The 2 groups analyzed (bilateral watershed infarcts and other infarct patterns) were not homogeneous at univariate analysis. As underscored by the authors, patients with bilateral watershed infarcts were more likely to have undergone an aortic procedure. Aortic procedures are at higher risk of neurological events because they involve wide aortic manipulation² which is considered a major risk factor for embolization.³ Patients with aortic dissections were found to have strong dominance of bilateral cerebral infarct, which was different from the unilateral pattern of "cardiac-type" operations.⁴ Moreover, surgery of the ascending aorta is generally performed with a different pattern of cardiopulmonary bypass as ascending aortic cannulation is avoided⁵ and involves more frequently deep hypothermic circulatory arrest.²

Even the duration of cardiopulmonary bypass was significantly different between the 2 groups. Cardiopulmonary bypass

machine produces systemic inflammatory response, destruction of blood cells, emboli, tissue edema and organ dysfunction.⁶ These effects lead to organ failure and poor outcomes if cardiopulmonary bypass time is increased.

In conclusion, the authors analyzed the effects of hypoperfusion on watershed strokes and gave new helpful information. However, the multifactorial bases of strokes still remain unclear and the relative role of hypoperfusion as well as embolization needs further study to be clarified.

Disclosures

None.

Fabio Barili, MD
Luca Dainese, MD
Gianluca Polvani, MD
Paolo Biglioli, MD

Department of Cardiovascular Surgery
Centro Cardiologico Monzino
University of Milan
Milan, Italy

1. Gottesman RF, Sherman PM, Grega MA, Yousem DM, Borowicz LM Jr, Selnes OA, Baumgartner WA, McKhann GM. Watershed strokes after cardiac surgery: diagnosis, etiology, and outcome. *Stroke*. 2006;37:2306–2311.
2. Goldstein LJ, Davies RR, Rizzo JA, Davila JJ, Cooperberg MR, Shaw RK, Kopf GS, Elefteriades JA. Stroke in surgery of the thoracic aorta: incidence, impact, etiology, and prevention. *J Thorac Cardiovasc Surg*. 2001;122:935–945.
3. Abu-Omar Y, Balacumaraswami L, Pigott DW, Matthews PM, Taggart DP. Solid and gaseous cerebral microembolization during off-pump, on-pump, and open cardiac surgery procedures. *J Thorac Cardiovasc Surg*. 2004;127:1759–1765.
4. Hedberg M, Boivie P, Edstrom C, Engstrom KG. Cerebrovascular accidents after cardiac surgery: an analysis of CT scans in relation to clinical symptoms. *Scand Cardiovasc J*. 2005;39:299–305.
5. Fusco DS, Shaw RK, Tranquilli M, Kopf GS, Elefteriades JA. Femoral cannulation is safe for type A dissection repair. *Ann Thorac Surg*. 2004;78:1285–1289.
6. Murphy GJ, Angelini GD. Side effects of cardiopulmonary bypass: what is the reality? *J Card Surg*. 2004;19:481–488.