

This Provisional PDF corresponds to the article as it appeared upon acceptance. Copyedited and fully formatted PDF and full text (HTML) versions will be made available soon.

17beta-estradiol effects on human coronaries and grafts employed in myocardial revascularization: a preliminary study

Journal of Cardiothoracic Surgery 2006, **1**:46 doi:10.1186/1749-8090-1-46

Gianluca Polvani (gianluca.polvani@unimi.it)
Fabio Barili (fabarili@libero.it)
Giuseppe Rossoni (giuseppe.rossoni@unimi.it)
Luca Dainese (luca.dainese@ccfm.it)
Manuela Wally Ossola (mossola@unimi.it)
Veli K Topkara (vk101@columbia.edu)
Francesco Grillo (francescogrill@ccfm.it)
Eleonora Penza (elepenza@unimi.it)
Elena Tremoli (elentrem@unimi.it)
Paolo Biglioli (paolo.biglioli@unimi.it)

ISSN 1749-8090

Article type Research article

Submission date 10 November 2006

Acceptance date 20 December 2006

Publication date 20 December 2006

Article URL <http://www.cardiothoracicsurgery.org/content/1/1/46>

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Journal of Cardiothoracic Surgery* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Journal of Cardiothoracic Surgery* or any BioMed Central journal, go to

<http://www.cardiothoracicsurgery.org/info/instructions/>

For information about other BioMed Central publications go to

© 2006 Polvani *et al.*, licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

<http://www.biomedcentral.com/>

© 2006 Polvani *et al.*, licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**17 β -estradiol effects on human coronaries and grafts employed in myocardial revascularization:
a preliminary study**

Gianluca Polvani¹, Fabio Barili¹, Giuseppe Rossoni², Luca Dainese¹, Manuela Wally Ossola¹, Veli K. Topkara³, Francesco Grillo¹, Eleonora Penza¹, Elena Tremoli¹, Paolo Biglioli¹.

¹*Department of Cardiovascular Surgery, University of Milan*

Centro Cardiologico Monzino

Via Parea 4, 20138 Milan, Italy

Phone +39 02 58002355 Fax +39 02 58011194

²*Department of Pharmacological Sciences, University of Milan*

Via Balzaretti 9, 20133 Milan, Italy

³*Division of Cardiothoracic Surgery, College of Physicians and Surgeon of Columbia University –
New York Presbyterian Hospital,*

Columbia University Medical Center, Milstein Hospital Building, 7GN-435

177 Fort Washington Avenue, New York, NY 10032, USA.

Corresponding Author:

Fabio Barili, MD

Department of Cardiovascular Surgery – University of Milan

Centro Cardiologico Monzino

Via Parea 4, 20138 Milan, Italy

Phone +39 02 58002355 Fax +39 02 580011194

E-mail: fabarili@libero.it fabio.barili@unimi.it

Abstract

Background. This study was undertaken to compare the *in vitro* effects of 17 β -estradiol on human epicardial coronary arteries, resistance coronary arteries and on arterial vessels usually employed as grafts in surgical myocardial revascularization.

Methods. Coronary artery rings (descending coronary artery, right coronary artery, circumflex coronary artery, first septal branch) and arterial graft rings (internal thoracic artery, gastro-epiploic artery) obtained from human heart donors with heart not suitable to cardiac transplantation were connected to force transducer for isometric force recording. Precontracted specimens with and without endothelium were exposed to increasing concentration of 17 β -estradiol (3-30-300-3000 nmol/l) and to vehicle (0.1% v/v ethanol). We also evaluated the effects of 17 β -estradiol on vessels before and 20 minutes after exposure to L-monomethyl-arginine and indomethacin.

Results. 17 β -estradiol induced a significant relaxation in all precontracted vessels (mean maximum effect: 78,6% \pm 8,5). This effect was not different among the different rings and was not related to the presence of endothelium. N-monomethyl-L-arginine and indomethacin did not modify 17 β -estradiol relaxant effect.

Conclusions. The vasodilator action of the 17 β -estradiol is similar on coronary arteries, resistance coronary arteries and arterial vessels usually employed as grafts in myocardial revascularization.

TEXT

Background

The interest for 17β -estradiol as vasoactive and vasoprotective agent is raising, since it was observed that it takes effect directly on the vascular wall, improving vasodilatation and inhibiting neointimal proliferation [1-7]. New devices, such as 17β -estradiol-eluting- stents, were developed to protect revascularized heart [8], hypothesizing a new role of 17β -estradiol for tertiary prevention in coronary artery disease (CAD).

Coronary perfusion after coronary artery bypass grafting (CABG) is a complex system dependent on several factors, including gender and the type of grafts employed [9-14]. Women have smaller coronary arteries than men and it can lead to incomplete revascularization and increased risk of in-hospital mortality [10]. The type of graft employed can affect outcomes as arterial grafts permit superior long-term patency and lower mortality rate [11-14]. Moreover, blood flow distribution in cardiac wall is related to not only diastolic pressure and section area of epicardial vessels but also depends on the resistance to the blood flow determined by intramyocardial branches, i.e. first septal branch [15].

The effect of estrogen on coronary system after surgical myocardial revascularization should be evaluated considering together all vessels that permit blood circulation, including grafts, epicardial coronary arteries and resistance vessels. To date, a comprehensive evaluation of estrogenic action on coronary arteries system was not performed. This study was undertaken to compare the effect of 17β -estradiol on epicardial coronary arteries, resistance coronary vessels and arteries employed as grafts in CABG.

Methods

We evaluated the *in vitro* effect of 17 β -estradiol on human epicardial coronary arteries (anterior interventricular artery, right coronary artery, circumflex artery) resistance vessels (first septal branch) and arteries usually employed in CABG (left internal mammary artery, gastroepiploic artery).

The vessels were obtained from 11 human donors whose heart was not suitable for cardiac transplantation and was harvested for banking cryopreserved valvular homografts. All patients were female (mean age 38 ± 11 years, range 18-54 years). All women had normal coronary arteries, without macroscopic atherosclerotic process.

Coronaries were dissected within 1 hours after the removal of heart and all segments were immediately put in a modified Krebs solution (composition in mmol/l: NaCl 118.3, KCl 4.7, CaCl₂ 2.5, MgSO₄ 1.2, KH₂PO₄ 1.2, NaHCO₃ 25, EDTA Calcium 0.026, glucose 11.1, albumin 0.1) at 4° C to be conserved. At the time of experiment, at maximum 1 hour after dissection, the vessels were cut into 3-4 mm long rings and suspended in an organ bath containing 10 ml modified Krebs solution aerated constantly with 95% O₂ and 5% CO₂ and maintained at 37° C. Each ring was mounted on a triangular-shaped metal hook connected to force transducer for isometric force recordings. Resting force was 2 grams [16].

Each single experiment was conducted on two rings from the same vessel. In one ring endothelium was mechanically removed with a wooden applicator (Group 1) in order to simulate an atheromatic vessel. The ring was precontracted with prostaglandin F_{2 α} (PGF_{2 α} , 1 μ mol/l), then histamine (0.1 μ mol/l) was added and the absence of vasodilative effect confirmed the complete removal of the endothelium (Figure 1-A). In the other ring from the same vessel, endothelium was preserved (Group 2). It was precontracted with prostaglandin F_{2 α} (PGF_{2 α} , 1 μ mol/l) and exposed to the same concentration of histamine (0.1 μ mol/l) with subsequent complete vasodilatation (Figure 2-B).

The rings (Group1 and Group2) were washed out and precontracted with prostaglandin F_{2 α} (PGF_{2 α} , 1 μ mol/l). After the contraction plateau was reached (about 10 minutes), rings were exposed to the vehicle (0.1% v/v ethanol, Figure 1-C). The bath solution was replaced, the rings were

precontracted again with prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$, 1 $\mu\text{mol/l}$) and, at the contraction plateau, they were exposed to increasing concentrations of 17β -estradiol (3-30-300-3000 nmol/l , Figure 1-D).

At the end of the experiment, the rings were washed out and we evaluated the effects of increasing concentrations of 17β -estradiol after the exposition to L-monomethyl-arginine (L-NMMA, 0.1 mmol/l) and indomethacin (10 $\mu\text{mol/l}$), L-NMMA is a non-specific inhibitor of nitric oxide synthase (NOS) that permits to evaluate the role of nitric oxide (NO) in vasoactive action of estrogen. Indomethacin is a cyclo-oxygenase inhibitor that blocks endothelial synthesis of prostacyclin. The rings were pretreated with both L-NMMA and indomethacin together for 20 minutes and precontracted with prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$, 1 $\mu\text{mol/l}$). After the contraction plateau was reached (about 10 minutes), rings were exposed to increasing concentrations of 17β -estradiol (3-30-300-3000 nmol/l).

This study had the approval of our Institutional Ethics Committee.

Statistical analysis.

The effect of 17β -estradiol on vessels was expressed as the percentage relaxation of the maximum contraction induced by $PGF_{2\alpha}$. Continuous variables were expressed as mean \pm standard deviation of the mean (SD). Differences between two groups were evaluated using Student's *t*-test. Repeated-measures analysis of variance (ANOVA) was used to compare more than two means. If statistically significant, Student's paired *t* test was then performed, with Bonferroni's method used to correct for multiple comparisons. A *p* value of less than 0.05 was considered statistically significant. Statistical analyses were performed using SPSS 13.0 software (SPSS, Inc, Chicago, IL).

Results

17 β -estradiol induced significant relaxation of precontracted coronary artery segments and vessels employed in CABG (compared with vehicle solvent, $p < 0.05$, data not shown). This vascular response to 17 β -estradiol was concentration-dependent with a maximum effect at 3 $\mu\text{mol/l}$ -dose (mean maximum effect: $78.6\% \pm 8.5\%$).

There were no significant differences in vasorelaxation between different types of vessels ($p > 0.05$, $n = 22$ in each group; Figure 2). It suggests that estrogen effect on vascular system is not dependent on vascular district and on segment's size.

The relaxant effect of estrogen was similar in groups with and without endothelium ($p > 0.05$, $n = 66$ in each group, Figure 3), suggesting an endothelium-independent mechanism of action. L-NMMA and indomethacin did not significantly inhibit the relaxation produced by increasing concentrations of 17 β -estradiol (Figure 4), excluding a role of nitric oxide and prostacyclin on estrogen-dependent relaxation.

Discussion

The relationship between 17β -estradiol and heart has been widely evaluated in the last decade, since it was observed that the risk of coronary artery disease significantly increases in women after menopause [17]. Several clinical studies focused on the protective role of postmenopausal HRT with contrasting results that leave the debate opened [18,19]. We shifted the attention on tertiary prevention of CAD to understand the vasoactive effects of 17β -estradiol on all conduits of the revascularized heart.

Our main question regarded the eventual diverse effects of estrogen on resistance vessel, epicardial vessels and arteries commonly employed as graft in CABG. Several studies evaluated only epicardial vessels [16, 20-23], without considering the importance of resistance vessels on heart perfusion. Moreover, LIMA graft was found responsible to estrogen but no comparison with coronaries was performed [24]. This study demonstrated a global acute vasorelaxant response of all vessels to 17β -estradiol which ameliorates all the complex physiology of blood flow in heart and arterial grafts. Hence, estrogen can acutely increase myocardial perfusion in women after coronary artery bypass grafting, acting through both a vasodilatation of coronary epicardial vessels and grafts and a decrease in resistance offered by resistance vessels.

The estrogenic vasoactive effect was found similar on normal and endothelium-deprived segments, confirming previous data on epicardial vessels [16, 22, 23]. Impaired-endothelium is characteristic of atherosclerotic coronaries and diseased arterial grafts. Surgical maneuvers are demonstrated to impair graft's endothelium and coronaries at incision site, worsening the endothelial function and leading to the well-known graft disease. The endothelium-independent vasorelaxation can be helpful in preventing perioperative vasoconstriction due to impaired endothelium and arterial graft spasm [23, 24]. Moreover, estrogens accelerate endothelial cells growth, increasing local expression of vascular endothelial growth factors and inhibiting endothelial cells apoptosis [25]. This estrogen-related rapid reendothelialization, as well as vasorelaxation and inhibition of neointimal proliferation, led to the development of new estrogen-eluting stents [8] and can also represent

protective effects for surgical revascularized heart. It could be useful especially in female sex, in which perioperative and postoperative complications are increased by an unfavorable anatomy [9, 10].

The vasorelaxant mechanisms of 17β -estradiol on vascular conduits are far to be completely clarified [26]. New data about non-genomic mechanisms of action lead to consider 17β -estradiol also as an acute and mid-term vasodilator. 17β -estradiol both stimulates endothelial NO production in a non-genomic manner and has vasorelaxant effects on impaired vessels acting on the muscular layer through an endothelium-independent mechanism. Smooth muscular cells respond to estrogens stimulating myocyte NO-synthesis or through similar Ca-antagonist mechanisms [16, 23, 27, 28]. Our study confirms the similar Ca-antagonist mechanism, as L-NMMA (N-monomethyl-L-arginine) does not change vascular response, even if we did not evaluate the myocyte NO-synthesis.

Limitations of the study

This study was performed on *in vitro* specimens and the concomitant *in vivo* effects could not be evaluated. By its nature, it did not considered chronic estrogenic effects that can be related to different mechanisms, such as genomic induction. Moreover, we focused the attention on endothelium-independent mechanisms similar to Ca-antagonist, as they are responsible of vasodilatation on both normal and impaired vessels, which are characteristic of revascularized heart.

Conclusions

This study demonstrated that 17β -estradiol has a similar relaxant effect on human female coronary arteries (epicardial capacitance arteries and resistance vessels) and arteries used as graft in CABG. Acute estrogenic administration can have vasorelaxant effect on all female revascularized heart, thus protecting coronaries and grafts and favoring reendothelialization.

List of abbreviations

ANOVA	analysis of variance
CABG	coronary artery bypass grafting
CAD	coronary artery disease
HRT	hormone replacement therapy
L-NMMA	L-monomethyl-arginine
LIMA	left internal mammary artery
NO	nitric oxide
NOS	nitric oxide synthase
PGF _{2α}	prostaglandin F _{2α}
SD	standard deviation of the mean

Competing interests

The authors declare that they have no competing interests'.

Authors' contribution

GP conceived of the study, and participated in its design and helped to draft the manuscript. **FB** participated in the study's design and coordination, performed the statistical analysis and drafted the manuscript. **LD** harvested hearts form human donors and harvested all specimens for the study. **EP** **FG** and **ET** participated in the study's design and helped to perform the statistical analysis. **VKT** participated in the study's design helped to draft the manuscript and edited it. **GR** and **MVO** carried out the "in vitro" experiments. **PB** coordinated the study and participated in its design.

All authors read and approved the final manuscript.

Acknowledgement

None.

References

1. Matsubara Y, Murata M, Kawano K, Zama T, Aoki N, Yoshino H, Watanabe G, Ishikawa K, Ikeda Y. Genotype distribution of estrogen receptor polymorphism in men and postmenopausal women from healthy and coronary populations and its relation to serum lipid levels. *Arterioscler Thromb Vasc Biol* 1997;17(11):3006-3012.
2. Venkov CD, Rankin AB, Vaughan DE. Identification of authentic estrogen receptor in cultured endothelial cells. A potential mechanism for steroid hormone regulation of endothelial function. *Circulation* 1996;94(4):727-733.
3. Karas RH, Patterson BL, Mendelsohn ME. Human vascular smooth muscle cells contain functional estrogen receptor. *Circulation* 1994;89(5):1943-1950.
4. Lin AL, Gonzales R, Carey KD, Shain SA. Estradiol-17 β affects estrogen receptor distribution and elevates progesterone receptor content in baboon aorta. *Arteriosclerosis* 1986;6:495-504.
5. Krasinski K, Spyridopoulos I, Asahara T, Van Der Zee R, Isner JM, Losordo DW. Estradiol accelerates functional endothelial recovery after arterial injury. *Circulation* 1997;95:1768-1772.
6. Chandrasekar B, Tanguay JF. Local delivery of 17-beta-estradiol decreases neointimal hyperplasia after coronary angioplasty in a porcine model. *J Am Coll Cardiol* 2000;36(6):1972-8.
7. Chandrasekar B, Nattel S, Tanguay JF. Coronary artery endothelial protection after local delivery of 17beta-estradiol during balloon angioplasty in a porcine model: a potential new pharmacologic approach to improve endothelial function. *J Am Coll Cardiol* 2001;38(5):1570-6.
8. Abizaid A, Albertal M, Costa MA, Abizaid AS, Staico R, Feres F, Mattos LA, Sousa AG, Moses J, Kipshidize N, Roubin GS, Mehran R, New G, Leon MB, Sousa JE. First human experience with the 17-beta-estradiol-eluting stent: the Estrogen And Stents To Eliminate Restenosis (EASTER) trial. *J Am Coll Cardiol* 2004;43(6):1118-21.
9. Corbineau H, Lebreton H, Langanay T, Logeais Y, Leguerrier A. Prospective evaluation of coronary arteries: influence on operative risk in coronary artery surgery. *Eur J Cardiothorac Surg* 1999;16(4):429-434.

10. O'Connor NJ, Morton JR, Birkmeyer JD, Olmstead EM, O'Connor GT. Effect of coronary artery diameter in patients undergoing coronary bypass surgery. *Circulation* 1996;93(4):652-655.
11. Motwani JG, Topol EJ. Aortocoronary saphenous vein graft disease: pathogenesis, predisposition, and prevention. *Circulation* 1998;97(9):916-931.
12. He GW. Arterial grafts for coronary surgery: vasospasm and patency rate. *J Thorac Cardiovasc Surg* 2001;121(3):431-433.
13. Boylan MJ, Lytle BW, Loop FD, Taylor PC, Borsh JA, Goormastic M, Cosgrove DM. Surgical treatment of isolated left anterior descending coronary stenosis. Comparison of left internal mammary artery and venous autograft at 18 to 20 years of follow up. *J Thorac Cardiovasc Surg* 1994;107(3):657-662.
14. Loop FD, Lytle BW, Cosgrove DM, Stewart RW, Goormastic M, Williams GW, Golding LA, Gill CC, Taylor PC, Sheldon WC, et al. Influence of the internal-mammary-artery graft on 10-years survival and other cardiac events. *N Engl J Med* 1986;314(1):1-6.
15. Chilian WM, Eastham CL, Layne SM, Marcus ML. Small vessel phenomena in the coronary microcirculation: phasic intramyocardial perfusion and coronary microvascular dynamics. *Progr Cardiovasc Dis* 1988;31(1):17-38.
16. Mugge A, Riedel M, Barton M, Kuhn M, Lichtlen PR. Endothelium independent relaxation of human coronary arteries by 17 β -oestradiol in vitro. *Cardiovasc Res* 1993;27(11):1939-1942.
17. Mendelsohn ME, Karas RH. The protective effect of estrogen on the cardiovascular system. *N Engl J Med* 1999;340(23):1801-1811.
18. Sullivan JM, El-Zeky F, Vander Zwaag R, Ramanathan KB. Effect on survival of estrogen replacement therapy after coronary artery bypass grafting. *Am J Cardiol* 1997;79(7):847-850.
19. Paoletti R, Wenger NK. Review of the international position paper on women's health and menopause. A comprehensive approach. *Circulation* 2003;107:1336-1339.
20. Jiang CW, Sarrel PM, Lindsay DC, Poole-Wilson PA, Collins P. Endothelium-independent relaxation of rabbit coronary artery by 17 β -estradiol in vitro. *Br J Pharmacol* 1991;104(4):1033-1037.
21. Williams JK, Adams MR, Klopfenstein HS. Estrogen modulates responses of atherosclerotic coronary arteries. *Circulation* 1990;81(5):1680-1687.

22. Reis SE, Gloth ST, Blumenthal RS, Resar JR, Zacur HA, Gerstenblith G, Brinker JA.. Ethinyl estradiol acutely attenuates abnormal coronary vasomotor responses to acetylcholine in postmenopausal women. *Circulation* 1994;89(1):52-60.
23. Chester AH, Jiang C, Borland JA, Yacoub MH, Collins P. Oestrogen relaxes human pericardial coronary arteries through non-endothelium-dependent mechanism. *Coron Artery Dis* 1995;6(5):417-422.
24. Polvani G, Marino MR, Roberto M, Dainese L, Parolari A, Pompilio G, Di Matteo S, Fumero A, Cannata A, Barili F, Biglioli P. Acute effects of 17 β -estradiol on left internal mammary graft after coronary artery bypass surgery. *Ann Thorac Surg* 2002;74:695-699.
25. Chandrasekar B, Sirois MG, Geoffroy P, Lauzier D, Nattel S, Tanguay JF. Local delivery of 17beta-estradiol improves reendothelialization and decreases inflammation after coronary stenting in a porcine model. *Thromb Haemost* 2005;94(5):1042-7.
26. Mendelsohn ME. Genomic and nongenomic effects of estrogen in vasculature. *Am J Cardiol* 2002;90(1A):3F-6F.
27. White RE, Han G, Maunz M, Dimitropoulou C, El-Mowafy AM, Barlow RS, Catravas JD, Snead C, Carrier GO, Zhu S, Yu X. Endothelium-independent effect of estrogen on Ca²⁺-activated K⁺ channels in human coronary artery smooth muscle cells. *Cardiovasc Res* 2002;53(3):650–661.
28. White RE, Darkow DJ, Lang JL. Estrogen relaxes coronary arteries by opening BKCa channels through a cGMP-dependent mechanism. *Circ Res* 1995;77:936-942.

Figure legends




Figure 1. A recording showing the relaxant effect of 17β -estradiol on human female coronary arteries. In panel A, histamine had no effects on a denuded ring. In panel B, endothelium was preserved and histamine had a relaxant effect. In panel C, a ring was precontracted with $\text{PGF}_{2\alpha}$ and exposed to the solvent (ethanol), without vasorelaxation. Adding $3\ \mu\text{M}$ of 17β -estradiol to the organ bath with ethanol, the force transducer recorded the maximum decrease in force within 10 minutes. Panel D shows the dose-dependent relaxation of a precontracted ring exposed to increasing concentrations of 17β -estradiol (3-30-300-3000 nmol/l).

Figure 2. The effects of 17β -estradiol on different vessels with and without endothelium. The effect of 17β -estradiol on vessels is expressed as the percentage relaxation of the maximum contraction induced by $\text{PGF}_{2\alpha}$. The relaxant effect of 17β -estradiol at each dose was similar in all groups ($n = 22$ for each group, $p > 0.05$ by repeated-measures analysis of variance). No intra-group significant difference was found between the same vessels with and without endothelium (data not shown). 17β -estradiol has a similar vasoactive effect on both epicardial coronaries and septal branch and arteries usually used as graft in myocardial revascularization at each concentration.


Figure 3. The effect of increasing concentrations (3-30-300-3000 nmol/l) of 17β -estradiol on vessels with and without endothelium. The relaxation is expressed as the percentage of the maximum effect obtained with $\text{PGF}_{2\alpha}$. The peak tension with $\text{PGF}_{2\alpha}$ was 3.2 ± 1.3 for intact vessels and 3.4 ± 1.0 for denuded vessels. There were no significant differences in vascular response to estrogen in groups with or without endothelium at each estrogen concentration ($n=66$ for group with endothelium, $n=66$ for group without endothelium, $p > 0.05$).

Figure 4. The effect of L-NMMA (0.1 mmol/l) and indomethacin (10 $\mu\text{mol/l}$) on 17β -estradiol vasorelaxation. The relaxation is expressed as the percentage of the maximum effect obtained with

PGF_{2α}. The peak tension with PGF_{2α} was 3.3±0.9 for experiment before exposure to L-NMMA and indomethacin and 3.1±1.2 for the experiments after exposure to L-NMMA and indomethacin. We did not find significant differences among groups at each 17β-estradiol concentration (n=132 in each group, p>0.05). No intra-group significant difference was found between different vessels and between vessels with and without endothelium (p>0.05, data not shown).



Relaxation
(% precontraction)




- ◇— Descending Coronary Artery
- x— Circumflex Artery
- Right Coronary Artery
- △- Internal Mammary Artery
- Gastroepiploic Artery
- *— First Septal Branch

17β-ESTRADIOL (nmol/l)

Figure 2

17 β -ESTRADIOL (nmol/l)



With endothelium
Without endothelium

17 β -ESTRADIOL (nmol/l)

